Liberia (DHS 2007)

Table - Coverage of Countdown indicators at national level and by key equity stratifiers.

	Family planning needs satisfied	Contraceptive prevalence rate	Antenatal care (1+ visits, skilled provider)	Antenatal care (4+ visits, any provider)	Skilled attendant at delivery	C-section rate	Early initiation of breastfeeding	Postnatal care for babies born at home	Postnatal care for all babies	BCG vaccine	DTP3 vaccine	Measles vaccine	Vitamin A in last 6 mos.	ORT and continued feeding	Care seeking for pneumonia	ITN use by children	Improved drinking water source	Co-coverage (6 or more interventions)	Composite coverage index
National	24.3	11.4	79.3	66.0	46.3	3.5	67.4	11.5	49.3	77.1	50.5	63.0	43.0	47.3	62.2	NO	66.2	11.2	51.0
Wealth quintiles																			
Q1	10.0	3.6	67.3	54.9	25.8	0.8	66.1	9.5	30.3	62.7	29.8	45.1	33.9	40.0	58.5	NO	28.3	5.1	37.5
Q2	13.3	5.7	69.2	56.7	31.9	2.0	72.1	10.4	35.4	64.8	38.2	55.2	36.7	40.9	45.8	NO	58.3	7.1	39.0
Q3	27.2	14.2	77.9	67.9	42.6	3.6	71.4	15.6	49.8	78.4	55.9	64.0	42.5	51.7	72.1	NO	75.4	12.3	54.1
Q4	28.9	15.4	91.8	76.3	68.7	5.7	62.8	11.5	64.4	94.1	66.0	74.4	52.6	52.6	62.4	NO	81.7	16.3	61.2
Q5	39.2	20.4	95.8	78.3	81.3	7.5	62.9	10.4	75.7	92.1	71.9	86.2	56.7	55.9	89.0	NO	87.3	20.9	70.3
Equity indicators																			
Q5/Q1 ratio	3.9	5.7	1.4	1.4	3.2	9.7	1.0	1.1	2.5	1.5	2.4	1.9	1.7	1.4	1.5	NO	3.1	4.1	1.9
Q5-Q1 difference	29.2	16.8	28.5	23.5	55.5	6.7	-3.2	1.0	45.4	29.4	42.1	41.1	22.8	15.9	30.5	NO	59.0	15.9	32.7
Concentration index	23.5	30.1	8.2	8.0	24.3	37.7	-1.7	8.6	19.5	9.5	17.1	12.2	11.2	6.7	9.0	NO	17.7	27.8	13.4
SII	35.0	22.1	38.5	32.3	63.0	8.6	-7.0	4.5	54.1	43.9	52.3	45.9	28.3	20.9	32.7	NO	66.1	19.2	43.8
Maternal education																			
none	19.1	7.7	74.2	61.5	35.7	1.9	69.6	9.5	39.7	71.5	46.8	57.9	40.6	48.7	56.5	NO	NA	8.1	46.2
primary	22.4	12.1	79.0	65.9	47.5	4.3	67.9	15.0	50.3	76.7	44.0	60.8	41.6	44.4	63.5	NO	NA	11.5	49.4
secondary +	37.8	20.9	92.8	77.1	75.7	6.4	60.6	9.9	71.6	90.1	70.3	78.4	53.3	49.5	77.2	NO	NA	20.4	65.8
Child's sex																			
male	NA	NA	77.7	67.4	46.6	3.6	68.0	11.7	50.5	74.5	49.2	61.3	43.6	48.4	63.6	NO	NA	11.8	NA
female	NA	NA	81.0	64.4	45.9	3.4	66.7	11.3	48.0	80.0	52.0	65.0	42.3	46.1	60.8	NO	NA	10.6	NA
Area of residence																			
urban	35.4	18.8	94.4	76.1	78.7	5.4	62.9	10.7	71.5	91.6	69.8	76.7	53.2	49.6	72.9	NO	83.7	18.9	65.5
rural	17.5	7.7	71.7	60.9	32.2	2.6	69.7	11.7	38.2	70.1	41.2	56.4	38.5	46.4	58.9	NO	55.8	8.0	44.0
Country region																			
monrovia	36.3	19.0	95.6	75.9	83.8	6.3	61.9	10.9	75.9	93.7	75.1	79.8	59.1	50.8	83.7	NO	83.8	21.3	68.8
north central	19.4	8.5	63.3	63.4	32.6	3.2	78.2	14.5	45.4	71.5	46.4	59.8	37.5	47.9	64.5	NO	60.7	9.1	45.6
north western	18.3	9.6	82.0	67.5	47.6	2.8	76.3	11.4	36.9	85.2	64.0	67.8	29.0	57.1	62.8	NO	71.8	10.1	52.7
south central	21.3	10.2	92.9	68.1	43.1	2.7	64.0	6.3	36.4	76.5	39.5	57.8	47.8	53.5	49.1	NO	51.4	9.1	48.6
south eastern a	22.9	9.7	79.1	51.3	32.8	2.0	42.6	8.3	37.0	68.1	28.0	51.7	27.6	37.9	38.4	NO	58.4	6.6	41.1
south eastern b	14.8	6.1	78.4	53.5	30.6	0.9	49.8	11.3	35.9	49.6	19.1	40.0	47.2	31.2	69.6	NO	56.7	5.0	38.9



Figure 1 - Coverage levels in the poorest and richest quintiles for selected interventions along the continuum of care.



Figure 2 - Coverage levels in the five wealth quintiles for selected interventions along the continuum of care.



Figure 3 - Co-coverage of health interventions: percentage of children aged 1–4 years according to the number of key child-survival interventions received, by wealth quintile.



Figure 4 - Composite coverage of selected interventions and corresponding coverage gap (how much is needed to reach universal coverage), by wealth quintile.