



Countdown to 2015

Maternal, Newborn & Child Survival

Country profiles

The information summarized in the profiles is intended to help policy makers and their partners assess progress, prioritize actions, and ensure accountability for national commitments to reduce maternal, newborn, and child mortality.

The following section contains profiles for 73 countries¹³ with high burdens of maternal and/or child mortality, including the 49 lowest-income countries covered by the Global Strategy for Women's and Children's Health. Profiles are included for the following countries:

Afghanistan	Gambia	Niger
Angola	Ghana	Nigeria
Azerbaijan	Guatemala	Pakistan
Bangladesh	Guinea	Papua New Guinea
Benin	Guinea-Bissau	Peru
Bolivia	Haiti	Philippines
Botswana	India	Rwanda
Brazil	Indonesia	Sao Tome and Principe
Burkina Faso	Iraq	Senegal
Burundi	Kenya	Sierra Leone
Cambodia	Korea, Democratic People's	Solomon Islands
Cameroon	Republic of	Somalia
Central African Republic	Kyrgyzstan	South Africa
Chad	Lao People's Democratic Republic	Swaziland
China	Lesotho	Tajikistan
Comoros	Liberia	Tanzania, United Republic of
Congo	Madagascar	Togo
Congo, Democratic Republic of the	Malawi	Turkmenistan
Côte d'Ivoire	Mali	Uganda
Djibouti	Mauritania	Uzbekistan
Egypt	Mexico	Vietnam
Equatorial Guinea	Morocco	Yemen
Eritrea	Mozambique	Zambia
Ethiopia	Myanmar	Zimbabwe
Gabon	Nepal	

13. There are 75 Countdown countries. Profiles for Sudan and South Sudan are not provided because of lack of available data at the time of publication.



Countdown to 2015

Maternal, Newborn & Child Survival

Gabon


March 2012

DEMOGRAPHICS

Total population (000)	1,505	(2010)
Total under-five population (000)	185	(2010)
Births (000)	41	(2010)
Birth registration (%)	89	(2000)
Total under-five deaths (000)	3	(2010)
Neonatal deaths: % of all under-5 deaths	37	(2010)
Neonatal mortality rate (per 1000 live births)	26	(2010)
Infant mortality rate (per 1000 live births)	54	(2010)
Stillbirth rate (per 1000 total births)	17	(2009)
Total maternal deaths	100	(2008)
Lifetime risk of maternal death (1 in N)	110	(2008)
Total fertility rate (per woman)	3.3	(2010)
Adolescent birth rate (per 1000 women)	144	(1998)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 55 (2010)

Percent HIV+ pregnant women receiving ARVs for PMTCT




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2000

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2000)	Early initiation of breastfeeding (within 1 hr of birth, %)	71	(2000)
Low birthweight incidence (moderate and severe, %)	14	(2000)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	0	(2009)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Gambia


March 2012

DEMOGRAPHICS

Total population (000)	1,728	(2010)
Total under-five population (000)	287	(2010)
Births (000)	66	(2010)
Birth registration (%)	55	(2005-2006)
Total under-five deaths (000)	6	(2010)
Neonatal deaths: % of all under-5 deaths	34	(2010)
Neonatal mortality rate (per 1000 live births)	31	(2010)
Infant mortality rate (per 1000 live births)	57	(2010)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	250	(2008)
Lifetime risk of maternal death (1 in N)	49	(2008)
Total fertility rate (per woman)	4.9	(2010)
Adolescent birth rate (per 1000 women)	104	(2000)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2005-2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2005-2006)
Low birthweight incidence (moderate and severe, %) 20 (2005-2006)

Early initiation of breastfeeding (within 1 hr of birth, %) 48 (2005-2006)
Introduction of solid, semi-solid/soft foods (%) -
Vitamin A two dose coverage (%) 100 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Ghana


March 2012

DEMOGRAPHICS

Total population (000)	24,392	(2010)
Total under-five population (000)	3,533	(2010)
Births (000)	770	(2010)
Birth registration (%)	71	(2008)
Total under-five deaths (000)	57	(2010)
Neonatal deaths: % of all under-5 deaths	38	(2010)
Neonatal mortality rate (per 1000 live births)	28	(2010)
Infant mortality rate (per 1000 live births)	50	(2010)
Stillbirth rate (per 1000 total births)	22	(2009)
Total maternal deaths	2,600	(2008)
Lifetime risk of maternal death (1 in N)	66	(2008)
Total fertility rate (per woman)	4.2	(2010)
Adolescent birth rate (per 1000 women)	70	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 0 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	9	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	52	(2008)
Low birthweight incidence (moderate and severe, %)	13	(2008)	Introduction of solid, semi-solid/soft foods (%)	76	(2008)
			Vitamin A two dose coverage (%)	93	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Guatemala


March 2012

DEMOGRAPHICS

Total population (000)	14,389	(2010)
Total under-five population (000)	2,167	(2010)
Births (000)	467	(2010)
Birth registration (%)	-	-
Total under-five deaths (000)	14	(2010)
Neonatal deaths: % of all under-5 deaths	48	(2010)
Neonatal mortality rate (per 1000 live births)	15	(2010)
Infant mortality rate (per 1000 live births)	25	(2010)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	480	(2008)
Lifetime risk of maternal death (1 in N)	210	(2008)
Total fertility rate (per woman)	4.0	(2010)
Adolescent birth rate (per 1000 women)	92	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 1998-1999

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	1	(2008-2009)
Low birthweight incidence (moderate and severe, %)	11	(2008-2009)
Early initiation of breastfeeding (within 1 hr of birth, %)	56	(2008-2009)
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	36	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Guinea


March 2012

DEMOGRAPHICS

Total population (000)	9,982	(2010)
Total under-five population (000)	1,658	(2010)
Births (000)	390	(2010)
Birth registration (%)	43	(2005)
Total under-five deaths (000)	48	(2010)
Neonatal deaths: % of all under-5 deaths	30	(2010)
Neonatal mortality rate (per 1000 live births)	38	(2010)
Infant mortality rate (per 1000 live births)	81	(2010)
Stillbirth rate (per 1000 total births)	24	(2009)
Total maternal deaths	2,700	(2008)
Lifetime risk of maternal death (1 in N)	26	(2008)
Total fertility rate (per woman)	5.2	(2010)
Adolescent birth rate (per 1000 women)	153	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 27 (2010)

Percent HIV+ pregnant women receiving ARVs for PMTCT




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2005

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	40	(2005)
Low birthweight incidence (moderate and severe, %)	12	(2005)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	97	(2010)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health




DEMOGRAPHICS

Total population (000)	1,515	(2010)
Total under-five population (000)	240	(2010)
Births (000)	58	(2010)
Birth registration (%)	24	(2010)
Total under-five deaths (000)	8	(2010)
Neonatal deaths: % of all under-5 deaths	28	(2010)
Neonatal mortality rate (per 1000 live births)	40	(2010)
Infant mortality rate (per 1000 live births)	92	(2010)
Stillbirth rate (per 1000 total births)	30	(2009)
Total maternal deaths	650	(2008)
Lifetime risk of maternal death (1 in N)	18	(2008)
Total fertility rate (per woman)	5.1	(2010)
Adolescent birth rate (per 1000 women)	137	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition


Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 9 (2010)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: Poorest 20% (red circles) Richest 20% (orange circles)




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	6	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	55	(2010)
Low birthweight incidence (moderate and severe, %)	11	(2010)	Introduction of solid, semi-solid/soft foods (%)	43	(2010)
			Vitamin A two dose coverage (%)	100	(2010)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	9,993	(2010)
Total under-five population (000)	1,237	(2010)
Births (000)	266	(2010)
Birth registration (%)	81	(2005-2006)
Total under-five deaths (000)	45	(2010)
Neonatal deaths: % of all under-5 deaths	16	(2010)
Neonatal mortality rate (per 1000 live births)	27	(2010)
Infant mortality rate (per 1000 live births)	70	(2010)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	820	(2008)
Lifetime risk of maternal death (1 in N)	93	(2008)
Total fertility rate (per woman)	3.3	(2010)
Adolescent birth rate (per 1000 women)	69	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 45 (2010)

Percent HIV+ pregnant women receiving ARVs for PMTCT




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2005-2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 10 (2005-2006)
Low birthweight incidence (moderate and severe, %) 25 (2005-2006)

Early initiation of breastfeeding (within 1 hr of birth, %) 44 (2005-2006)
Introduction of solid, semi-solid/soft foods (%) 90 (2005-2006)
Vitamin A two dose coverage (%) 21 (2010)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	1,224,614	(2010)
Total under-five population (000)	127,979	(2010)
Births (000)	27,165	(2010)
Birth registration (%)	41	(2005-2006)
Total under-five deaths (000)	1,696	(2010)
Neonatal deaths: % of all under-5 deaths	52	(2010)
Neonatal mortality rate (per 1000 live births)	32	(2010)
Infant mortality rate (per 1000 live births)	48	(2010)
Stillbirth rate (per 1000 total births)	22	(2009)
Total maternal deaths	63,000	(2008)
Lifetime risk of maternal death (1 in N)	140	(2008)
Total fertility rate (per woman)	2.6	(2010)
Adolescent birth rate (per 1000 women)	45	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




NFHS 2005-2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	20	(2005-2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	41	(2007-2008)
Low birthweight incidence (moderate and severe, %)	28	(2005-2006)	Introduction of solid, semi-solid/soft foods (%)	56	(2005-2006)
			Vitamin A two dose coverage (%)	34	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Indonesia


March 2012

DEMOGRAPHICS

Total population (000)	2,39,871	(2010)
Total under-five population (000)	21,579	(2010)
Births (000)	4,372	(2010)
Birth registration (%)	53	(2007)
Total under-five deaths (000)	151	(2010)
Neonatal deaths: % of all under-5 deaths	48	(2010)
Neonatal mortality rate (per 1000 live births)	17	(2010)
Infant mortality rate (per 1000 live births)	27	(2010)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	10,000	(2008)
Lifetime risk of maternal death (1 in N)	190	(2008)
Total fertility rate (per woman)	2.1	(2010)
Adolescent birth rate (per 1000 women)	52	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition


Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 100 (2010)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	15	(2007)	Early initiation of breastfeeding (within 1 hr of birth, %)	44	(2007)
Low birthweight incidence (moderate and severe, %)	9	(2007)	Introduction of solid, semi-solid/soft foods (%)	85	(2007)
			Vitamin A two dose coverage (%)	80	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health




DEMOGRAPHICS

Total population (000)	31,672	(2010)
Total under-five population (000)	5,188	(2010)
Births (000)	1,125	(2010)
Birth registration (%)	95	(2006)
Total under-five deaths (000)	43	(2010)
Neonatal deaths: % of all under-5 deaths	53	(2010)
Neonatal mortality rate (per 1000 live births)	20	(2010)
Infant mortality rate (per 1000 live births)	31	(2010)
Stillbirth rate (per 1000 total births)	9	(2009)
Total maternal deaths	710	(2008)
Lifetime risk of maternal death (1 in N)	300	(2008)
Total fertility rate (per woman)	4.7	(2010)
Adolescent birth rate (per 1000 women)	68	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 0 (2010)

Percent HIV+ pregnant women receiving ARVs for PMTCT

Uncertainty range around the estimate


No Data

Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	6	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	31	(2006)
Low birthweight incidence (moderate and severe, %)	15	(2006)	Introduction of solid, semi-solid/soft foods (%)	62	(2006)
			Vitamin A two dose coverage (%)	-	

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	40,513	(2010)
Total under-five population (000)	6,664	(2010)
Births (000)	1,529	(2010)
Birth registration (%)	60	(2008-2009)
Total under-five deaths (000)	122	(2010)
Neonatal deaths: % of all under-5 deaths	35	(2010)
Neonatal mortality rate (per 1000 live births)	28	(2010)
Infant mortality rate (per 1000 live births)	55	(2010)
Stillbirth rate (per 1000 total births)	22	(2009)
Total maternal deaths	7,900	(2008)
Lifetime risk of maternal death (1 in N)	38	(2008)
Total fertility rate (per woman)	4.7	(2010)
Adolescent birth rate (per 1000 women)	106	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) ²⁰¹⁰

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008-2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) ⁷ (2008-2009)
Low birthweight incidence (moderate and severe, %) ⁸ (2008-2009)
Early initiation of breastfeeding (within 1 hr of birth, %) ⁵⁸ (2008-2009)
Introduction of solid, semi-solid/soft foods (%) ⁸⁵ (2008-2009)
Vitamin A two dose coverage (%) ⁶² (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	24,346	(2010)
Total under-five population (000)	1,704	(2010)
Births (000)	348	(2010)
Birth registration (%)	100	(2009)
Total under-five deaths (000)	12	(2010)
Neonatal deaths: % of all under-5 deaths	52	(2010)
Neonatal mortality rate (per 1000 live births)	18	(2010)
Infant mortality rate (per 1000 live births)	26	(2010)
Stillbirth rate (per 1000 total births)	13	(2009)
Total maternal deaths	810	(2008)
Lifetime risk of maternal death (1 in N)	230	(2008)
Total fertility rate (per woman)	2.0	(2010)
Adolescent birth rate (per 1000 women)	1	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Bigible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	5	(2009)	Early initiation of breastfeeding (within 1 hr of birth, %)	18	(2009)
Low birthweight incidence (moderate and severe, %)	6	(2009)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	99	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Kyrgyzstan


March 2012

DEMOGRAPHICS

Total population (000)	5,334	(2010)
Total under-five population (000)	595	(2010)
Births (000)	130	(2010)
Birth registration (%)	94	(2006)
Total under-five deaths (000)	5	(2010)
Neonatal deaths: % of all under-5 deaths	46	(2010)
Neonatal mortality rate (per 1000 live births)	19	(2010)
Infant mortality rate (per 1000 live births)	33	(2010)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	97	(2008)
Lifetime risk of maternal death (1 in N)	450	(2008)
Total fertility rate (per woman)	2.7	(2010)
Adolescent birth rate (per 1000 women)	30	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 19 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	65	(2006)
Low birthweight incidence (moderate and severe, %)	5	(2006)	Introduction of solid, semi-solid/soft foods (%)	60	(2006)
			Vitamin A two dose coverage (%)	97	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health




DEMOGRAPHICS

Total population (000)	6,201	(2010)
Total under-five population (000)	683	(2010)
Births (000)	141	(2010)
Birth registration (%)	72	(2006)
Total under-five deaths (000)	8	(2010)
Neonatal deaths: % of all under-5 deaths	39	(2010)
Neonatal mortality rate (per 1000 live births)	21	(2010)
Infant mortality rate (per 1000 live births)	42	(2010)
Stillbirth rate (per 1000 total births)	14	(2009)
Total maternal deaths	980	(2008)
Lifetime risk of maternal death (1 in N)	49	(2008)
Total fertility rate (per woman)	2.7	(2010)
Adolescent birth rate (per 1000 women)	110	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 85 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	30	(2006)
Low birthweight incidence (moderate and severe, %)	11	(2006)	Introduction of solid, semi-solid/soft foods (%)	41	(2006)
			Vitamin A two dose coverage (%)	83	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	2,171	(2010)
Total under-five population (000)	274	(2010)
Births (000)	60	(2010)
Birth registration (%)	45	(2009)
Total under-five deaths (000)	5	(2010)
Neonatal deaths: % of all under-5 deaths	42	(2010)
Neonatal mortality rate (per 1000 live births)	35	(2010)
Infant mortality rate (per 1000 live births)	65	(2010)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	310	(2008)
Lifetime risk of maternal death (1 in N)	62	(2008)
Total fertility rate (per woman)	3.2	(2010)
Adolescent birth rate (per 1000 women)	92	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 24 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2009)	Early initiation of breastfeeding (within 1 hr of birth, %)	53	(2009)
Low birthweight incidence (moderate and severe, %)	13	(2004)	Introduction of solid, semi-solid/soft foods (%)	68	(2009)
			Vitamin A two dose coverage (%)	38	(2007)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	3,994	(2010)
Total under-five population (000)	681	(2010)
Births (000)	154	(2010)
Birth registration (%)	4	(2007)
Total under-five deaths (000)	15	(2010)
Neonatal deaths: % of all under-5 deaths	34	(2010)
Neonatal mortality rate (per 1000 live births)	34	(2010)
Infant mortality rate (per 1000 live births)	74	(2010)
Stillbirth rate (per 1000 total births)	27	(2009)
Total maternal deaths	1,400	(2008)
Lifetime risk of maternal death (1 in N)	20	(2008)
Total fertility rate (per woman)	5.2	(2010)
Adolescent birth rate (per 1000 women)	177	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 18 (2010)

Percent HIV+ pregnant women receiving ARVs for PMTCT




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	44	(2010)
Low birthweight incidence (moderate and severe, %)	14	(2007)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	97	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	20,714	(2010)
Total under-five population (000)	3,305	(2010)
Births (000)	732	(2010)
Birth registration (%)	80	(2008-2009)
Total under-five deaths (000)	44	(2010)
Neonatal deaths: % of all under-5 deaths	37	(2010)
Neonatal mortality rate (per 1000 live births)	22	(2010)
Infant mortality rate (per 1000 live births)	43	(2010)
Stillbirth rate (per 1000 total births)	21	(2009)
Total maternal deaths	3,000	(2008)
Lifetime risk of maternal death (1 in N)	45	(2008)
Total fertility rate (per woman)	4.7	(2010)
Adolescent birth rate (per 1000 women)	147	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 6 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008-2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	15	(2003-2004)	Early initiation of breastfeeding (within 1 hr of birth, %)	72	(2008-2009)
Low birthweight incidence (moderate and severe, %)	16	(2008-2009)	Introduction of solid, semi-solid/soft foods (%)	86	(2008-2009)
			Vitamin A two dose coverage (%)	95	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Malawi


March 2012

DEMOGRAPHICS

Total population (000)	14,901	(2010)
Total under-five population (000)	2,715	(2010)
Births (000)	663	(2010)
Birth registration (%)	-	-
Total under-five deaths (000)	56	(2010)
Neonatal deaths: % of all under-5 deaths	32	(2010)
Neonatal mortality rate (per 1000 live births)	27	(2010)
Infant mortality rate (per 1000 live births)	58	(2010)
Stillbirth rate (per 1000 total births)	24	(2009)
Total maternal deaths	3,000	(2008)
Lifetime risk of maternal death (1 in N)	36	(2008)
Total fertility rate (per woman)	6.0	(2010)
Adolescent birth rate (per 1000 women)	177	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 26 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	58	(2006)
Low birthweight incidence (moderate and severe, %)	13	(2006)	Introduction of solid, semi-solid/soft foods (%)	86	(2010)
			Vitamin A two dose coverage (%)	96	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Mali


March 2012

DEMOGRAPHICS

Total population (000)	15,370	(2010)
Total under-five population (000)	2,912	(2010)
Births (000)	714	(2010)
Birth registration (%)	81	(2010)
Total under-five deaths (000)	120	(2010)
Neonatal deaths: % of all under-5 deaths	28	(2010)
Neonatal mortality rate (per 1000 live births)	48	(2010)
Infant mortality rate (per 1000 live births)	99	(2010)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	4,500	(2008)
Lifetime risk of maternal death (1 in N)	22	(2008)
Total fertility rate (per woman)	6.3	(2010)
Adolescent birth rate (per 1000 women)	190	(2004)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 40 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	9	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	46	(2006)
Low birthweight incidence (moderate and severe, %)	19	(2006)	Introduction of solid, semi-solid/soft foods (%)	25	(2006)
			Vitamin A two dose coverage (%)	99	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Mauritania


March 2012

DEMOGRAPHICS

Total population (000)	3,460	(2010)
Total under-five population (000)	513	(2010)
Births (000)	117	(2010)
Birth registration (%)	56	(2007)
Total under-five deaths (000)	13	(2010)
Neonatal deaths: % of all under-5 deaths	36	(2010)
Neonatal mortality rate (per 1000 live births)	39	(2010)
Infant mortality rate (per 1000 live births)	75	(2010)
Stillbirth rate (per 1000 total births)	27	(2009)
Total maternal deaths	590	(2008)
Lifetime risk of maternal death (1 in N)	41	(2008)
Total fertility rate (per woman)	4.5	(2010)
Adolescent birth rate (per 1000 women)	88	(2002)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 54 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	7	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	81	(2010)
Low birthweight incidence (moderate and severe, %)	34	(2007)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	97	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Mexico


March 2012

DEMOGRAPHICS

Total population (000)	113,423	(2010)
Total under-five population (000)	11,095	(2010)
Births (000)	2,217	(2010)
Birth registration (%)	-	-
Total under-five deaths (000)	37	(2010)
Neonatal deaths: % of all under-5 deaths	44	(2010)
Neonatal mortality rate (per 1000 live births)	7	(2010)
Infant mortality rate (per 1000 live births)	14	(2010)
Stillbirth rate (per 1000 total births)	5	(2009)
Total maternal deaths	1,700	(2008)
Lifetime risk of maternal death (1 in N)	500	(2008)
Total fertility rate (per woman)	2.3	(2010)
Adolescent birth rate (per 1000 women)	90	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics


No Data

NUTRITION

Wasting prevalence (moderate and severe, %)	2	(2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	18	(2009)
Low birthweight incidence (moderate and severe, %)	7	(2009)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	-	-

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Morocco


March 2012

DEMOGRAPHICS

Total population (000)	31,951	(2010)
Total under-five population (000)	3,022	(2010)
Births (000)	623	(2010)
Birth registration (%)	85	(2000)
Total under-five deaths (000)	23	(2010)
Neonatal deaths: % of all under-5 deaths	53	(2010)
Neonatal mortality rate (per 1000 live births)	19	(2010)
Infant mortality rate (per 1000 live births)	30	(2010)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	720	(2008)
Lifetime risk of maternal death (1 in N)	360	(2008)
Total fertility rate (per woman)	2.3	(2010)
Adolescent birth rate (per 1000 women)	18	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2003-2004

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization

— Percent of children immunized against measles
— Percent of children immunized with 3 doses DTP
— Percent of children immunized with 3 doses Hib




Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider

■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	11 (2003-2004)	Early initiation of breastfeeding (within 1 hr of birth, %)	52 (2003-2004)
Low birthweight incidence (moderate and severe, %)	15 (2003-2004)	Introduction of solid, semi-solid/soft foods (%)	-
		Vitamin A two dose coverage (%)	43 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health




DEMOGRAPHICS

Total population (000)	23,391	(2010)
Total under-five population (000)	3,876	(2010)
Births (000)	883	(2010)
Birth registration (%)	31	(2008)
Total under-five deaths (000)	114	(2010)
Neonatal deaths: % of all under-5 deaths	30	(2010)
Neonatal mortality rate (per 1000 live births)	39	(2010)
Infant mortality rate (per 1000 live births)	92	(2010)
Stillbirth rate (per 1000 total births)	28	(2009)
Total maternal deaths	4,800	(2008)
Lifetime risk of maternal death (1 in N)	37	(2008)
Total fertility rate (per woman)	4.9	(2010)
Adolescent birth rate (per 1000 women)	185	(2001)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) ⁹ (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2003

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	4	(2008)	Early initiation of breastfeeding (within 1 hr of birth, %)	63	(2008)
Low birthweight incidence (moderate and severe, %)	16	(2008)	Introduction of solid, semi-solid/soft foods (%)	86	(2008)
			Vitamin A two dose coverage (%)	100	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed






DEMOGRAPHICS

Total population (000)	47,963	(2010)
Total under-five population (000)	3,956	(2010)
Births (000)	830	(2010)
Birth registration (%)	72	(2009-2010)
Total under-five deaths (000)	56	(2010)
Neonatal deaths: % of all under-5 deaths	47	(2010)
Neonatal mortality rate (per 1000 live births)	32	(2010)
Infant mortality rate (per 1000 live births)	50	(2010)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	2,400	(2008)
Lifetime risk of maternal death (1 in N)	180	(2008)
Total fertility rate (per woman)	2.0	(2010)
Adolescent birth rate (per 1000 women)	17	(2001)

Under-five mortality rate


Deaths per 1000 live births



Source: IGMIE 2011

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 28 (2010)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
— Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	8	(2009-2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	76	(2009-2010)
Low birthweight incidence (moderate and severe, %)	9	(2009-2010)	Introduction of solid, semi-solid/soft foods (%)	-	-
			Vitamin A two dose coverage (%)	94	(2010)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed

